

## 10.5 Cont'd

Ex: Find the area of one petal  
of  $r = \sin 3\theta$


Solve  $r = 0$  to find where petals start.

$$\sin 3\theta = 0$$

$$3\theta = 0, \pi, 2\pi, \dots$$


$$\theta = 0, \frac{\pi}{3}, \frac{2\pi}{3}, \dots$$

Integrate over  $0 \leq \theta \leq \frac{\pi}{3}$ .



$$\begin{aligned} A &= \frac{1}{2} \int_{\alpha}^{\beta} [f(\theta)]^2 d\theta \\ &= \frac{1}{2} \int_0^{\frac{\pi}{3}} \sin^2 3\theta d\theta \\ &= \frac{1}{2} \int_0^{\frac{\pi}{3}} \left[ \frac{1}{2} - \frac{\cos 6\theta}{2} \right] d\theta \\ &= \frac{1}{2} \left[ \frac{\theta}{2} - \frac{\sin 6\theta}{12} \right]_0^{\frac{\pi}{3}} \\ &= \frac{1}{2} \left[ \frac{\pi}{6} - 0 \right] \\ &= \frac{\pi}{12} \end{aligned}$$

Ex: Find the area inside  $r = 3 + 2\sin\theta$   
and outside  $r = 2$ .



Intersection :  $r = r$

$$3 + 2\sin\theta = 2$$

$$1 = -2\sin\theta$$

$$-\frac{1}{2} = \sin\theta$$

$$\theta = \frac{7\pi}{6}, \frac{11\pi}{6}$$

$$\text{or } \theta = \frac{7\pi}{6}, -\frac{\pi}{6}$$

Caution:  $\theta$  must be increasing

$$-\frac{\pi}{6} \leq \theta \leq \frac{7\pi}{6}$$

$$A = A_1 - A_2$$

area inside  
 $r = 3 + 2\sin\theta$

area inside  
 $r = 2$

$$A_1 = \frac{1}{2} \int_{-\frac{\pi}{6}}^{\frac{7\pi}{6}} (3 + 2\sin\theta)^2 d\theta$$

$$= \frac{1}{2} \int_{-\frac{\pi}{6}}^{\frac{7\pi}{6}} (9 + 12\sin\theta + \underbrace{4\sin^2\theta}_{2 - 2\cos 2\theta}) d\theta$$

$$= \frac{1}{2} \int_{-\frac{\pi}{6}}^{\frac{7\pi}{6}} (11 + 12\sin\theta - 2\cos 2\theta) d\theta$$

$$= \frac{1}{2} \left[ 11\theta - 12\cos\theta - \sin 2\theta \right]_{-\frac{\pi}{6}}^{\frac{7\pi}{6}}$$

$$= \frac{1}{2} \left[ \frac{77\pi}{6} - 12\left(-\frac{\sqrt{3}}{2}\right) - \frac{\sqrt{3}}{2} - \left(-\frac{11\pi}{6} - 12\left(\frac{\sqrt{3}}{2}\right) + \frac{\sqrt{3}}{2}\right) \right]$$

$$= \frac{1}{2} \left[ \frac{88\pi}{6} + 22\frac{\sqrt{3}}{2} \right]$$

$$= \frac{22\pi}{3} + \frac{11\sqrt{3}}{2}$$

$$A_2 = \frac{1}{2} \int_{-\pi/6}^{7\pi/6} 4 d\theta$$

$$= \frac{1}{2} [4\theta]_{-\pi/6}^{7\pi/6}$$


$$= 2 \left[ \frac{7\pi}{6} + \frac{\pi}{6} \right]$$

$$= \frac{8\pi}{3}$$

$$A = A_1 - A_2$$

$$= \frac{14\pi}{3} + \frac{11\sqrt{3}}{2}$$

Ex: Find the area inside both  
 $r = 1 + \sin\theta$  and  $r = 3\sin\theta$ .



By symmetry, the area is double  
the area in  $0 \leq \theta \leq \frac{\pi}{2}$ .


Intersection :  $r = r$

$$3 \sin \theta = 1 + \sin \theta$$

$$2 \sin \theta = 1$$

$$\sin \theta = \frac{1}{2}$$

$$\theta = \frac{\pi}{6}, \frac{5\pi}{6}$$



On  $0 \leq \theta \leq \frac{\pi}{6}$ ,  $r = 3 \sin \theta$

$\frac{\pi}{6} \leq \theta \leq \frac{\pi}{2}$ ,  $r = 1 + \sin \theta$

$$A = \underbrace{\int_0^{\pi/6} [3 \sin \theta]^2 d\theta}_{A_1} + \underbrace{\int_{\pi/6}^{\pi/2} (1 + \sin \theta)^2 d\theta}_{A_2}$$

$$A_1 = \int_0^{\pi/6} 9 \sin^2 \theta d\theta$$

$$= \frac{9}{2} \int_0^{\pi/6} (1 - \cos 2\theta) d\theta$$

$$= \frac{3\pi}{4} - \frac{9\sqrt{3}}{8}$$

$$A_2 = \int_{\pi/6}^{\pi/2} \left( 1 + 2\sin\theta + \frac{1 - \cos 2\theta}{2} \right) d\theta$$

$$= \int_{\pi/6}^{\pi/2} \left[ \frac{3}{2} + 2\sin\theta - \frac{\cos 2\theta}{2} \right] d\theta$$

$$= \left[ \frac{3}{2}\theta - 2\cos\theta - \frac{\sin 2\theta}{4} \right]_{\pi/6}^{\pi/2}$$

$$= \frac{3\pi}{4} - \left( \frac{3\pi}{12} - 2\frac{\sqrt{3}}{2} - \frac{1}{4}\frac{\sqrt{3}}{2} \right)$$

$$= \frac{3\pi}{4} - \frac{\pi}{4} + \sqrt{3} + \frac{\sqrt{3}}{8}$$

$$= \frac{\pi}{2} + \frac{9\sqrt{3}}{8}$$

$$A = A_1 + A_2$$

$$= \frac{3\pi}{4} - \frac{9\sqrt{3}}{8} + \frac{\pi}{2} + \frac{9\sqrt{3}}{8}$$

$$= \frac{5\pi}{4}$$